



Published in final edited form as:

J Subst Abuse Treat. 2010 April ; 38(3): 263–274. doi:10.1016/j.jsat.2010.01.002.

A Randomized Pilot Study of the Engaging Moms Program for Family Drug Court

Gayle A. Dakof, (Ph.D)^a, Jeri B. Cohen, (J.D.)^b, Craig E. Henderson, (Ph.D)^c, Eliette Duarte, (M.A)^b, Maya Boustani, (M.S)^a, Audra Blackburn^a, Ellen Venzer, (J.D.)^b, and Sam Hawes, (M.S.)^c

^aCenter for Treatment Research on Adolescent Drug Abuse, Department of Epidemiology and Public Health, University of Miami Miller School of Medicine, Miami, Florida 33136 USA

^bJuvenile Division, 11th Judicial Circuit, Miami Dade County Florida USA

^cSam Houston State University, Huntsville, Texas, USA

Abstract

In response to the need for effective drug court interventions, the effectiveness of the Engaging Moms Program (EMP) versus intensive case management services (ICMS) on multiple outcomes for mothers enrolled in family drug court was investigated. In this intent-to-treat study, mothers ($N = 62$) were randomly assigned to either usual drug court care or the Engaging Moms drug court program. Mothers were assessed at intake, and 3, 6, 12, and 18 months following intake. Results indicated that at 18 months post drug court enrollment, 77% of mothers assigned to EMP versus 55% of mothers assigned to ICMS had positive child welfare dispositions. There were statistically significant time effects for both intervention groups on multiple outcomes including substance use, mental health, parenting practices, and family functioning. EMP showed equal or better improvement than ICMS on all outcomes. The results suggest that EMP in family drug court is a viable and promising intervention approach to reduce maternal addiction and child maltreatment.

Keywords

child maltreatment; drug courts; women; addiction

1. Introduction

The devastating effects of drug abuse on the lives of addicted parents, their children, and extended families are undeniable: family disruption, child maltreatment, poor developmental outcomes, and the continuation of multigenerational poverty and addiction (Grogan-Kaylor, Ruffolo, Ortega, & Clarke, 2008; Keller, Catalano, Haggerty, & Fleming, 2002; Kelley & Fals-Stewart, 2004; Stanger, Higgins, Bickel, Elk, Grabowski, Schmitz et al., 1999; Walsh, MacMillan & Jamieson, 2003). Given the association between parental drug use and child maltreatment (Walsh et al, 2003), it is not surprising that many drug-dependent parents,

© 2010 Elsevier Inc. All rights reserved.

Corresponding author: Gayle Dakof: 1120 NW 14th Street, Suite 1010, Miami, FL 33136. Tel: 305-243-3656. Fax: 305-243-3651. GDakof@med.miami.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

particularly mothers, are involved in the child welfare system. Between 50 and 80% of children involved in the child welfare system have a drug-dependent parent (Barth, Courtney, Duerr Berrick, & Albert, 1994; Curtis & McCulloch, 1993; Locke & Newcomb, 2004), and these families are characterized by particularly poor outcomes, including increased re-referral to child welfare (Wolock & Magura, 1996), placement instability (Connell, Bergeron, Katz, Saunders, & Tebes, 2007), increased number of out of home placements (Barth et al., 1994; Dore, Doris, & Wright, 1995), and termination of parental rights (Marcenko, Kemp, & Larson, 2000). Clearly, the family environment of children of substance abusing parents is often compromised by instability, neglect, and poor parenting practices leading some to suggest that, "...the mission of child welfare depends in large part on the success in providing services to address parental substance abuse" (Choi & Ryan, 2006, p. 313).

There is evidence to suggest that improving the functioning of the biological parents—especially reducing parental drug use—not only reduces the risk to their children (Catalano, Gainey, Fleming, Haggerty, & Johnson, 1999; Drapela & Mosher, 2007; Kelley & Fals-Stewart, 2008) but also improves child welfare outcomes (Green, Rockhill, & Furrer, 2007; Grella, Needell, Shi, & Hser, 2009). However, substance abuse treatment completion among parents who come in contact with the child welfare system is low (Choi & Ryan, 2006; Gregoire & Schultz, 2001).

Looking for solutions to the problem of parental substance use and child maltreatment, many communities have turned to drug courts (Harrell & Goodman, 1999; Huddleston, Marlowe, & Casebolt, 2008; Semidei, Radel, & Nolan, 2001). Adapted from the adult drug court model, family drug courts were established to enhance the effectiveness of child welfare agencies by increasing enrollment and retention in substance abuse treatment, motivating parents to address their addiction, and coordinating the many social services needed to stabilize families. Ultimately these courts aim to help parents "become emotionally, financially, and personally self-sufficient and to develop parenting and coping skills adequate for serving as an effective parent on a day-to-day basis" (p. 5, OJP, 1998). It should be recognized, however, that although family drug courts are based on the adult court model, there are features of family drug court that distinguish it from adult drug court, namely these courts do not operate in the criminal justice system, the majority of participants are female, and the court addresses dual issues of parental addiction and recovery as well as child safety and permanency (Edwards & Ray, 2005; Green, Furrer, Worcel, Burrus, & Finigan, 2009). Given these complexities, it has been argued that the mission of dependency drug court is more complex than the mission of adult criminal or juvenile delinquency drug courts (Bryan & Havens, 2008; Pach, 2009).

Although community and judicial acceptance of, and enthusiasm for, drug courts are broad and deep, there are few investigations of their effectiveness generally (Belenko, 2001; Wilson, Mitchell & MacKenzie, 2006), and even fewer on family drug courts in particular (Green, Furrer, Worcel, Burrus, & Finigan, 2009). A small number of evaluations of dependency drug courts indicate that drug court has promise (Boles, Young, Moore & DiPirro-Beard, 2007; Green, Furrer, Worcel, Burrus & Finigan, 2007; Haack, Alemi, Nemes, & Cohen, 2004; Worcel 2008, Green et al, 2009). However, it should be noted that each evaluation has serious limitations including, among others, non-randomized designs and samples that are lacking in minority representation.

There appears to be considerable variation in the structure, components, and practices over the more than 250 dependency drug courts in the United States (BJA, Drug Court Clearinghouse, 2009), and no consensus beyond identifying key general components. Most dependency drug courts share certain basic features including a non-adversarial relationship among the parties, comprehensive assessment of service needs, frequent court hearings and drug testing, intensive judicial supervision, enrollment in substance abuse treatment and other necessary services, and

court administered rewards and sanctions. In order to graduate from dependency drug courts, participants must have successfully completed substance abuse treatment, have a specified period of continuous abstinence, show evidence of a safe and stable living situation, spend a substantial period of time adequately performing the parent role, and have a life plan initiated and in place (e.g., employment, education, vocational training).

Most family drug courts employ court counselors whose job is to refer clients to substance abuse treatment and other court-ordered services, develop a recovery service plan, and monitor and report clients' ongoing progress to the court (Edwards & Ray, 2005). Although the influential roles of the drug court judge and the court-affiliated substance abuse treatment program on drug court outcomes has been noted (e.g., Edwards & Ray, 2005; NADAP, 1997; Henggeler, 2007), little attention has focused on the intervention models drug court counselors employ, nor on the quality of their work. Undoubtedly court counselors have a significant role in drug court; therefore, it seems reasonable to suggest that the effectiveness of drug court is at least partially due to the services provided by the drug court caseworkers.

The Engaging Moms Program (EMP), which is the focus of the current study, was initially developed as a family-oriented intervention aimed at facilitating treatment entry and retention among mothers of substance-exposed infants. In the first study of this approach (Dakof, Quille, Tejada, Alberga, Bandstra, & Szapocznik, 2003), mothers were recruited either from a public hospital where either the mother or their newborn tested positive for cocaine, or from child welfare after an infant abuse/neglect involving maternal cocaine use reported. All mothers were referred to an agency where they received an in-home psychosocial evaluation, and a substance abuse treatment referral. Mothers who agreed to participate in this study were then randomized to receive either community services as usual (i.e., scheduled intake appointment at treatment program; reminder phone calls before the scheduled appointment; follow-up phone calls following the appointment; and if necessary repeating the sequence two more times) or the EMP. The results showed that EMP successfully facilitated substance abuse treatment entry and initial retention of non-treatment seeking drug abusing mothers into drug treatment. Significantly more mothers assigned to EMP enrolled into substance abuse treatment than did mothers assigned to community services as usual (88% versus 46%), and 67% of the participants in EMP received at least four weeks of treatment compared with 38% of the mothers assigned to the usual services condition.

Given the promising results of the EMP with addicted mothers of infants, the growth of family drug courts throughout the nation, the lack of specific guidelines or intervention manuals for drug court counselors, and disappointing outcomes of our local dependency drug court, we adapted the EMP for use in a drug court context. We conducted a natural experiment on 80 consecutive enrollments: 37 who participated in the dependency drug court prior to the implementation of the EMP and 43 who received drug court with EMP. The results were very encouraging: 72% of the mothers receiving EMP versus 38% receiving case management services graduated from drug court, and at 15 months post-entry 70% of EMP mothers were reunified with their children versus a 40% reunification rate among mothers who received drug court with traditional case management services (Dakof, Cohen & Duarte, 2009). Although the results were encouraging, this study had numerous limitations (non-randomized design, judicial bias, limited outcome information), and therefore, was a hypothesis-generating and not a hypothesis-testing study.

This work, however, brought us to the current study, which is a true pilot test of the Engaging Moms for Family drug court intervention. The current study was designed to examine the effectiveness of EMP in comparison to Intensive Case Management Services (ICMS), on the primary goal of family drug courts—producing positive child welfare outcomes. A secondary,

but also important goal of the EMP was to support recovery from substance use and enhance family functioning among drug court participants.

2. Materials and methods

2.1. Participants

This study was implemented at the State of Florida 11th Circuit Judicial Juvenile Court in Miami, Florida. All mothers accepted into the family drug court, called Dependency Drug Court (DDC), were eligible for the study. DDC eligibility criteria were that parents had to be: (1) 18 years or older, (2) with at least one child adjudicated dependent; (3) have a diagnosis of substance abuse or dependence; (4) have a potential for family reunification (parents with severe cognitive, emotional or physical disorders, or who have had their parental right terminated previously were considered ineligible for re-unification); and (5) after consultation with their attorney, voluntarily enroll in drug court.

2.2 Procedures

The University of Miami Institutional Review Board approved all materials and procedures. Mothers who participated in this study were adjudicated in a single drug court with one judge (not the founding family drug court judge) presiding, and received the same types of substance abuse treatment, parenting interventions, and other services as ordered by the judge. The judge was blind to study hypotheses and aims. The only difference between the two conditions was the intervention administered by the drug court counselors: EMP versus ICMS. Upon enrollment in drug court, mothers were referred to the study. Study staff met with mothers to describe the study and obtain written informed consent prior to the first assessment session. It was emphasized to the mothers that participation in the study was voluntary and did not influence their status in drug court or their child welfare case generally, and that they had the right to discontinue participation in the study at any time. Mothers who refused to participate in the study were still able to participate in drug court and received standard drug court intensive case management services. After the baseline assessment, mothers were randomly assigned to either ICMS ($n = 31$) or EMP ($n = 31$). An urn randomization program was used to ensure equivalence on 4 key variables: age, ethnicity, number of children, and years using drugs. In this intent-to-treat design, all mothers were assessed at the following time points regardless of the extent of their participation in drug court: intake into the study, and 3, 6, 9, 12, and 18 months following intake. Participants were paid to complete the research assessment interviews: \$100 for the intake interview, \$50 for the 3, 6, and 9 month interviews, and \$100 for the 12 and 18 month interviews. All research data (e.g., questionnaires, research urine screens) were collected by University of Miami research staff who were not affiliated with the court, and records were kept confidential as required by the University of Miami IRB.

2.3 Drug Court Counselors

Five counselors, all female, participated in this study: 3 delivering EMP and 2 delivering ICMS. Counselors in each condition received 30 hours of initial training as well as ongoing weekly supervision in their respective approaches. Counselors in each condition received similar amounts of training and supervision in order to facilitate adherence to and quality control of each approach.

2.4 Interventions: Intensive Case Management Services and Engaging Moms Program

Mothers were expected to follow the same basic requirements of participating in family drug court (i.e., completing drug treatment and remaining drug free; completing parenting classes and demonstrating adequate parenting skills; participating in educational/vocational training,

domestic violence or other counseling as ordered by the judge; obtaining adequate and stable housing; and being gainfully employed or in school).

The dependency drug court program was a 12 – 15 month program organized into four phases. Progression through the phases was related to the mother's level of substance abuse treatment and compliance with court orders. Mothers were drug tested (urine screens) at each court hearing and in their substance abuse treatment programs. During the first month of drug court, mothers were required to attend weekly drug court hearings. Thereafter, if reports to the court indicated that the mother was progressing well, court hearings were typically reduced to twice monthly. During the second phase of the program, which lasted 3 months, clients continued to attend twice monthly hearings. In the third phase, which lasted another 3 months, the frequency of hearings was reduced to once per month. In the fourth and final phase, which extended to graduation from the drug court program, clients attended hearings every 6 to 12 weeks. This multi-phased process included a collaborative team approach that involved court counselors, child welfare workers, treatment providers, parent educators, and other social and health care service providers, as needed. Drug court counselors had contact with their clients, either in-person or on the phone, on a weekly basis through Phase 2, reducing to bi-weekly in Phase 3, and monthly in Phase 4. Workers were available more frequently on an as-needed basis. The caseload for drug court counselors was between 10 – 15 active cases. The only difference between the two groups (EMP and ICMS) was how the drug court counselors worked with the mothers. All other aspects of the programs were the same.

ICMS was closely aligned with the drug court case management services recommended by the National Drug Court Institute, the ICMS model provided five key case management functions: assessment, planning, linkage, monitoring, and advocacy within the context of a strong case manager-client therapeutic alliance (Monchick, Scheyett, & Pfeifer, 2006). The overall objective was to assess needs, engage in collaborative intervention planning, provide referral to suitable drug abuse treatment and other services, coordinate the system of care providing services to the mother, closely supervise and monitor compliance with court orders, advocate for the mother with service providers, and provide emotional support. Case managers in this system served as a liaison between the court, substance abuse treatment providers, child welfare, and the client. The case manager was responsible for referrals to treatment and other court-ordered services, developing a recovery service plan, monitoring and reporting clients' ongoing progress to the court, reducing any barriers to the delivery of treatment and other services, and providing emotional and practical support to the mother.

EMP is based on the theory and method of Multidimensional Family Therapy (Liddle, Dakof, & Diamond, 1991) and was adapted for use in family drug court. EMP was designed to help mothers succeed in drug court by complying with all court orders such as attending and benefiting from substance abuse and other intervention programs (e.g., domestic violence counseling, parenting classes), attending court sessions, remaining drug free, and demonstrating capacity to parent her children. EMP counselors conducted individual and conjoint sessions with the mother and her family, focusing on six core areas of change: (1) mother's motivation and commitment to succeed in drug court and to change her life, (2) the emotional attachment between the mother and her children, (3) relationships between the mother and her family of origin, (4) parenting skills, (5) mother's romantic relationships, and (6) emotional regulation, problem solving, and communication skills.

EMP counselors achieve change in the six core areas by conducting a series of integrated individual and family sessions (e.g. individual sessions with mother, individual sessions with family/partner, family and couple sessions). The intervention is organized in 3 stages: Stage 1: Alliance and Motivation, Stage 2: Behavioral Change, and Stage 3: Launch To an Independent Life.

In Stage 1, the counselor is focused on two goals: (1) building a strong therapeutic alliance with the mother and her family, and (2) enhancing mother and family motivation to participate in drug court and to change. EMP counselors provide support to both the mother and her family. They empower and validate; highlight strengths and competence; build confidence in the program; and are very compassionate, loving and nurturing. To enhance motivation, the EMP counselor highlights the pain, guilt and shame that the mother and her family have experienced, and the high stakes involved (e.g., losing child to the child welfare system), while simultaneously creating positive expectations and hope.

Stage 2 is focused on behavioral change in both the mother and her family/spouse. EMP has several goals for this stage. First, counselors enhance the emotional attachment between the mother and her children by working individually with the mother to help her explore her maternal role. Mother and children sessions designed to enhance the mother's commitment to her children are also provided. Equally important is enhancement of the attachment between the mother and her family of origin and/or spouse. This is accomplished by helping the family restrain from negativity and offer instrumental and emotional support to the mother. Considerable attention is devoted to repairing the mother's relationship with her family, which frequently has been damaged by past hurts, betrayals and resentments. Romantic relationships, typically with men, have often been a source of pain and distress for many of the mothers involved in the child welfare system. Hence, the EMP program addresses these relationships by helping the mother conduct a relationship life review, including examining tensions between having a romantic relationship and being a mother. The counselors help the mother examine the choices she has made, and continues to make in terms of romantic relationships, and teaches her how to make better decisions for herself and her children. EMP counselors also help the mother deal with slips, mistakes, setbacks and relapses in a non-punitive and therapeutic manner (i.e. forward looking). Finally, in Stage 2, the EMP specialist facilitates the mother's relationship with court personnel (judge, child welfare workers, and attorneys) and treatment or other service providers. The EMP counselor conducts "shuttle diplomacy" between the mother and service providers to prevent and resolve problems, and help the mother take full advantage of the services being provided to her. With respect to the court, the drug court counselors facilitate therapeutic jurisprudence in the courtroom by preparing mothers for court appearances and advocating for the mother in front of the judge and at weekly drug court case reviews.

In the final launching phase (Stage 3), the EMP counselor helps the mother prepare for an independent life by developing a practical and workable routine for everyday life; addressing how the mother will balance self care, children and work; outlining a plan to address common emergencies with children and families; and addressing how the mother will deal with potential problems, mistakes, slips, and relapses.

2.5 Measures

All data, with the exception of child welfare outcomes, which were gathered from court records, were gathered in face-to-face interviews with study participants at intake, and at 3, 6, 9, 12 and 18 months following intake. Because drug court was designed to last between 12 and 15 months, the 18-month follow-up is post-drug court participation. Measures described below were administered at all assessment points (except participant characteristics, which were measured at intake and child welfare status which was measured at 18 months). Trained research assistants, blind to intervention assignment and study hypotheses, administered measures in face-to-face interviews with the mothers. All the measures used in this study demonstrated adequate reliability and validity.

2.5.1. Background Information—Background questions gathered directly from the mothers included age of mother, race/ethnicity, number of children, educational attainment, income, marital status, drug of choice, age at first drug use, age at birth of first child, lifetime arrests, lifetime physical abuse, and lifetime sexual abuse.

2.5.2. Child Welfare Status—Information on child welfare status at 18 months was extracted from court records. Child welfare status was defined as follows: (1) sole custody with one or more children, (2) joint custody with one or more children, (3) permanent guardianship with relative without termination of mother's parental rights, (4) permanent guardianship with relative with termination of mother's parental rights, or (5) child placed in foster care with termination of mother's parental rights.

2.5.3. Addiction Severity Index—The Addiction Severity Index (ASI, 5th Ed.; McLellan, Alterman, Cacciola, Metzger, & O'Brien, 1992), is a semi structured interview that assesses the participant's past and current functioning in seven problem areas: medical status, employment/financial status, legal status, drug use, alcohol use, family/social relationships, and psychiatric problems. It is among the most widely used instruments in the field of addictions and has shown excellent reliability and validity (e.g. Carise, McLellan, Cacciola, Love, Cook, et al., 2001). Trained research assistants at all assessment points administered the ASI, and composite scores which indicate the present status of the mother were calculated. In this study, the drug use, alcohol use, mental health, employment, medical, and family/social relationship composite scores were used in the analyses reported below.

2.5.4 Urinalysis—Urine specimens were collected at all research assessment points, using the OnTrak TesTcup Pro8, an in vitro diagnostic test used for the qualitative detection of drug or drug metabolites in participant's urine. This method allowed for the identification of the presence of eight substances: alcohol, THC, amphetamines, barbiturates, benzodiazepines, cocaine, opiates, and PCP.

2.5.5. Revised Conflict Tactics Scale—The Revised Conflict Tactics Scale (R-CTS) is based on the CTS, developed over 39 years ago to assess family conflict and violence (Strauss, 1979). The CTS is widely used in the area of child maltreatment and has excellent psychometric properties. The Child Parent CTS physical violence scale was used in the current study ($\alpha = .73$).

2.5.6. Brief Child Abuse Potential I—The Brief Child Abuse Potential (B-CAP) is a brief version of the Child Abuse Potential Inventory (CAPI) designed to assess primarily physical child abuse. The B-CAP has seven scales including the 24-item risk scale that was used in the present study. The B-CAP has demonstrated adequate predictive validity (Ondersma, Chaffin, Simpson, & LeBreton, 2005). Cronbach's alpha for the current study was .79.

2.5.7. Brief Symptom Inventory (BSI)—The BSI is a 53-item psychological self-report symptom scale used to measure nine primary symptom dimensions and three global indices (Derogatis, 1993). The BSI is a shortened version of the Symptom Checklist-90, a widely used scale assessing current psychological distress and symptoms in both patient and non-patient populations. The BSI is highly correlated with the SCL-90, and also has demonstrated high reliability and validity in hundreds of studies (Derogatis & Savitz, 1999). In the present study, we used the Global Severity Index (GSI), which measures the level of overall psychological distress ($\alpha = .97$).

2.5.8. Stages of Change Readiness and Treatment Eagerness Scale—The Stages of Change Readiness and Treatment Eagerness Scale (SOCRATES) is designed to measure

client readiness to change substance use related behaviors. The 19-item version used in the present study (Miller & Tonigan, 1996) measures three factors of readiness to change: Taking Steps ($\alpha = .87$), Recognition ($\alpha = .80$), and Ambivalence ($\alpha = .92$). This is one of the most widely used instruments for measuring motivation to change among adults with substance abuse problems.

2.5.9. Working Alliance Inventory—The Working Alliance Inventory (WAI; Horvath & Symonds, 1991; Kivlighan & Shaughnessy, 1995) assesses the three components of therapeutic alliance: bonds ($\alpha = .85$), goals ($\alpha = .91$), and tasks ($\alpha = .82$).

2.5.10. Services Satisfaction Scale-16—The Service Satisfaction Scale-16 (SSS-16) (Greenfield & Attkisson, 1989) is designed to measure several components of satisfaction with outpatient services. It consists of five subscales (Manner and Skill, Perceived Outcome, Procedures, Accessibility, Waiting), and a total satisfaction score derived from all items ($\alpha = .99$). The SSS has been widely used, including with substance abusers (Attkisson & Greenfield, 1994; Parr & Greenfield, 2000). In the current study we adapted it for use in drug court, and participants were instructed to report on their satisfaction with drug court services. The total satisfaction score was used in the present study.

3. Results

3.1 Data Analytic Approach

The study examined the: (a) effects of drug court over time (through 18 months after intake), and (b) comparative effects of two drug court interventions. This comparison was conducted using a 2 (intervention condition) by 6 (time) repeated measures intent-to-treat design. The focus of family drug court is to use the principles of therapeutic jurisprudence to work with drug-addicted mothers to maintain their parental rights, possibly even reunifying them with their children, while ensuring their children's safety. Therefore, the primary outcome of family drug court is child welfare outcome, and we have highlighted this first. However, issues such as reducing parental substance use and improving parenting practices, mental and physical health status are critical to achieving positive child welfare outcomes, and we examined these issues in our secondary outcome analyses.

We conducted analyses in the following outcomes: (1) child welfare outcomes, (2) substance use (measured by the Addiction Severity Index [ASI] and urine assays), (3) family functioning (measured by the ASI), (4) parenting practices (measured by the Brief Child Abuse Potential and Conflict Tactics Scales), and (5) maternal mental and physical health (measured by the BSI and ASI). We were also interested in process variables such as the mother's satisfaction with drug court services (SSS), her motivation to change (SOCRATES) and alliance with her drug court counselor (WAI).

For most outcomes (the exceptions being alliance and motivation to change for reasons we describe below), we analyzed individual client change with the latent growth curve (LGC) modeling method (Duncan, Duncan, Strycker, Li, & Alpert, 1999). Although sometimes considered a large-sample technique, it can be successfully used with smaller samples (Liddle et al., 2009) and can result in greater power than traditional methods such as repeated measures analysis of variance (Delucchi & Bostrom, 1999). Missing data, which was rare (97% of follow-up assessments were completed), was handled with full information maximum likelihood (FIML) estimation, under the assumption that the data were missing at random (Little & Rubin, 1987).

Latent growth curve (LGC) modeling conducted with Mplus (Version 5.2; Muthén & Muthén, 1998–2009) proceeded in three stages. First, we tested a series of growth curve models,

representing possible forms of growth (e.g., no change, linear change, discontinuous change), to determine the overall shape of the individual change trajectories. Given the shape of the observed average outcome trajectories, we initially tested a piecewise growth model (Crawford, Pentz, Chou, Li, & Dwyer, 2003) with two distinct phases of growth representing change in the initial intervention period (between intake and the 3-month follow-up) and maintenance of initial gains between the 3-month and 18-month follow-up assessment. In order for this model to be identified, we constrained the variance of the initial growth phase to 0. Second, we added intervention condition to the models to test the impact of intervention type on initial status and change over time (i.e., the intercept and slope growth parameters). Intervention effects were demonstrated by a statistically significant slope parameter, as tested by the pseudo- z test—calculated by dividing the coefficient by its standard error—associated with treatment condition. Finally, for those outcomes that showed significant improvements across intervention condition, we entered graduation status (yes or no) as a covariate in the LGC models to examine the extent to which change in outcome was associated with whether mothers successfully graduated from drug court (or not). We used Full Information Maximum Likelihood estimation, under the assumption that the data were missing at random (Little & Rubin, 1987), to accommodate missing data.

We considered process variables—specifically mothers' alliances with their drug court specialists¹, motivation to change, and satisfaction with services—to be more static rather than dynamic factors (i.e., we considered these variables to be important factors at certain critical time points rather than outcome variables that we expected to change over time). Therefore, we used more conventional statistical methods (independent samples t -tests) to compare the two intervention conditions on these variables. In addition, we only report intervention comparisons, and not time effects, for these variables.

In line with recommendations of researchers for intervention development, our primary goal was to develop an acceptable, feasible, and reasonably promising intervention that could be subsequently tested in a larger randomized controlled trial (RCT; Rounsaville, Carroll, & Onken, 2001). Also consistent with these researchers' recommendations given the work that goes into initial intervention development (e.g., specifying the theoretical rationale and theory of change of the disorder, demonstrating feasibility of delivering the intervention, specifying process measures, developing a manual, testing the intervention on pre-pilot cases, etc.), the sample size was fairly small, and in some cases may be underpowered for testing intervention effects. Therefore, we report both significance tests and effect sizes (Cohen's d) associated with the intervention effects. Effect sizes were calculated using Feingold's (2009) method for growth curve modeling. We base our interpretations of intervention effects on effect sizes of $d = .5$ or larger (considered a medium effect size), assuming that subsequent RCTs would include sample sizes sufficient for detecting a medium effect size with statistical power of .80 or greater².

3.2 Baseline demographic and clinical characteristics

Participants were 62 mothers recruited from family drug court. Table 1 summarizes participant characteristics. There were no significant differences between conditions on any variables listed in Table 1. Mothers who participated in this study, were on average 30.2 years old ($SD = 11$), and were Black (42%), Hispanic (35%), White Non-Hispanic (23%). They had low incomes, reporting a median monthly family income of \$561, and 71% were unemployed at intake. Mothers were not well educated; only 43% had at least a high school school diploma

¹We examined change in working alliance using RM-ANOVA, and the results suggested that the alliances did not change over time.

²We conducted a post-hoc power analysis using Monte Carlo simulation methods and parameter estimates from the current study. The results indicated that a sample size of 140 participants was sufficient to produce power of .80 assuming a moderate-sized treatment effect of $d = .50$.

or GED. They had an average of 2.5 children, and only 10% reported being married. Many mothers were themselves victims of child maltreatment with 55% reporting a history of child physical abuse and 36% a history of child sexual abuse. They were primarily polydrug users, either with (48%) or without alcohol (19%). Sixteen percent identified cocaine as their drug of choice. Mothers had considerable mental health problems with 68% showing symptoms of serious depression, 55% with serious anxiety symptoms, and 19% reported current suicidal ideation. Finally, mothers reported that they began using drugs between the ages of 16 – 17, and averaged approximately 3 lifetime arrests.

3.3. Response and Attrition Rates

During the study period, 69 mothers were enrolled in drug court, 62 of whom enrolled in the study and 7 who declined, resulting in an 89% response rate. Attrition rates after randomization by assessment points were as follows: 3 months: 6%; 6 months: 6%; 9 months: 12%; 12 months: 8%; and 18 months 8%. Attrition rates did not differ by treatment condition. There were no missing administrative record data on child welfare status at 18 months.

3.4 Adherence

Similar intervention adherence monitoring procedures were followed for both conditions throughout the study. Supervisors, who were experts in either ICMS or EMP, provided close supervision to all court case managers. Supervisors for each intervention reviewed all active cases during weekly group supervision lasting approximately 2.5 hours. In order to demonstrate that counselors adhered to the basic parameters of the intervention (i.e., session frequency and duration, participants), drug court specialists completed contact logs for every contact with clients. As shown in Table 2, mothers in both intervention groups received a similar amount of contact from their court counselors. ICMS mothers received an average of 37.26 hours of service ($SD = 19.15$, Median = 42.45), and mothers in the EMP condition received an average of 38.48 hours of service ($SD = 20.26$, Median = 36.62). The two groups differed with respect to the amount of family sessions received with mothers in the EMP receiving significantly more hours of family session than mothers in ICMS (6.92 versus 3.83). This is consistent with the EMP model, which makes a concerted effort to include the family.

3.5. Child Welfare Status

Sixty-six percent of mothers had positive permanency outcomes, which we defined as sole custody, joint custody, or permanent guardianship with family members without termination of the mother's parental rights. As shown in table 3, participants randomized to EMP were more likely than mothers in ICMS to have positive child welfare outcomes 18 months after enrollment in drug court. Specifically, 77% of EMP mothers versus 55% of ICMS mothers had positive child welfare outcomes ($\chi^2 [1, N = 62] = 3.53, p = .060$).

3.6 Maternal Substance Use, Psychosocial, and Family Functioning

Significant improvement from intake through the 3-month follow-up was demonstrated on maternal substance use and family functioning, and these initial improvements were maintained through the 18-month follow-up (see tables 4 and 5). As shown in table 5, Mothers showed fairly steep declines in their self-reported alcohol and drug use, as well as their likelihood for testing positive for substance use on urine screens. These initial gains were maintained through 18 months, as demonstrated by non-significant growth in the remaining follow-up period.

Mothers also showed significant improvement in other domains of psychosocial functioning (mental health, medical problems, and employment). Between intake and 3 months, mothers showed fairly strong improvement on both the BSI Global Severity Index and the ASI

Psychiatric Status. Further, on the BSI, the mothers continued to improve between 3 and 18 months. Mothers did not show similar improvement on the ASI; however, they maintained their initial improvements (Mean Slope = -0.01 , $SE = 0.01$). Likewise, mothers showed improvement in their medical problems between intake and three months, and marginally significant improvement between 3 and 18 months. With respect to employment, after showing an initial increase in employment problems, they showed significant improvement between 3 and 18 months.

Regarding family functioning and parenting practices, mothers significantly improved their family functioning between intake and 3 months³ as well as decreasing their risk for child abuse. In the time period between 3 and 18 months, mothers continued to improve their family functioning and potential for child abuse.

When we compare the two intervention conditions, we found no statistically significant intervention effects; however, several outcomes favoring EMP showed effect sizes of $d = .5$ or greater. Please note that the effect sizes reported here are for the first 3 months of the follow-up period. In each case, these initial treatment differences were maintained through 18 months. As compared to women receiving ICMS, women receiving EMP were more likely to decrease their alcohol use ($d = 1.45$), experience improvement in their mental health (as measured by the BSI, $d = .50$), improve their overall family functioning ($d = .63$) and decrease their risk for child abuse ($d = .51$). Differences between the two interventions were negligible for drug use with mothers in both groups showing significant improvement over time. Regarding medical problems, although both treatments showed approximately equivalent change between intake and 3 months, those receiving EMP showed greater improvements between 3 and 18 months than those receiving ICMS.

3.7 Therapeutic Alliance, Motivation to Change, and Satisfaction with Services

With respect to process variables of therapeutic alliance, motivation to change and satisfaction with services received, mothers receiving EMP reported higher alliances with their drug court specialists at the 3-month follow-up (Tasks: $t [55] = 2.37$, $p = .021$; Bonds: $t [55] = 2.29$, $p = .026$; Goals: $t [55] = 2.14$, $p = .037$). There were no differences between conditions on motivation to change or satisfaction with services.

3.8 Drug court graduation in relation to substance use

We then examined associations between change in outcome variables and graduation status. As expected, mothers who successfully graduated from drug court showed steeper declines in substance use from 3 to 18 months than those who did not graduate (Mean Slope = -0.01 , $SE = 0.01$, $pseudo\ z = -2.08$, $p = .04$). Overall, 61% of the mothers graduated from drug court; 67% of EMP and 53% of ICMS mothers. This difference was not statistically significant.

4. Discussion

The results of this pilot study suggest that Engaging Moms Program shows promise in the family drug court context. Engaging Moms Program delivered in the drug court context increased the likelihood of positive outcomes in comparison to intensive case management. In all domains of functioning, families assigned to EMP showed improvement that was equal to or better than families assigned to ICMS. In particular the results suggest that the Engaging Moms in Family Drug Court intervention has significant promise with respect to child welfare outcomes, alcohol use, family functioning, risk for child abuse, and the mother's mental and physical health status. Although this study does not allow a clear explanation for why, for

³Note that given the way the ASI items are worded, decreases in the ASI Family/Social Status reflect improvements in family functioning.

example, mothers who received EMP had better child welfare outcomes than mothers who received ICMS, the data do suggest some possibilities. One could argue that the primary difference between the two models concerns therapeutic alliance and family involvement. Mothers participating in EMP reported significantly greater alliance with their drug court counselor than did mothers in ICMS, and this was true on all 3 subscales of a therapeutic alliance (task, bonds and goals). Next, EMP had almost twice as many family sessions as did ICMS, and thus it would not be unreasonable to speculate that involving the mother's family (i.e., parents, siblings, spouse/partner) facilitates success in family drug court. Although EMP involves more than developing a strong therapeutic alliance and involving the mother's family, these are essential components of the EMP. Developing a strong therapeutic alliance with the mother involves providing emotional support; highlighting her strengths and competencies; showing respect, empathy, and compassion; and generally empowering the mother, and as a result sets the foundation for subsequent work. Involvement of the mother's family, especially her family of origin, is obviously a key component of the family-based EMP. With respect to the family, the EMP first locates and reaches out to all family members who are important to the mother and may be able to help her recovery. We facilitate the family's involvement in the mother's life. Then the counselor conducts a series of specialized sessions designed to help the mother and family address and then move beyond any past hurts and betrayals that are often a part of the addict's family life. Finally, in a very practical way we encourage the family to provide both emotional and, if possible, practical social support to the mother and her children. It is not unreasonable to speculate that this kind of support can be the difference between reunification and termination of parental rights. Sometimes when a mother cannot be reunified with her children, if the family is involved in a positive way with the mother and her children, termination of parental rights can still be avoided. Many times the family takes over custody of the children without termination of the mother's rights. This allows the mother to be involved with her children and family even though she may not be able to be the custodial parent. In the EMP, then, we (a) get the mother's family involved in her life, (b) encourage the family to provide both practical and emotional support to the mother and her children, and (c) if the mother is not able to be reunified with her children, we encourage mother and family to come to an agreement whereby a family member is the primary parent but the mother is still involved with her children and family in a positive way. This is a key aspect of the Engaging Moms Program, and might be a reason for its' success in preventing termination of parental rights and maintaining family integrity.

Although the results of this pilot study are encouraging, there are important limitations. The primary limitation is that it has a small sample size that limits statistical power for testing intervention group differences, and increases the unreliability of any estimates of within-group or between-group effects due to error variance. It is possible that different results would be obtained with a larger sample. Although the reported effect sizes were in the medium to strong range, we were not able to adequately test for the statistical significance of intervention effects. An increased sample size may have uncovered more reliable and stable effects in the targeted domains. Second, there was no comparison of mothers in a non-drug court setting. Although the time effects are strong and significant, without an appropriate comparison the results cannot address whether or not drug court outcomes are better than non-drug court (i.e., ordinary dependency court) outcomes with drug abusing mothers.

Despite these limitations, this study also has some significant strengths. First, we were able to implement a fully randomized clinical trial in a family drug court setting. To our knowledge, this is the first study of its kind. Second, study methods are state-of the science. This study was an intent-to-treat longitudinal design with a high participant response rate (89%), a judge who was blind to study hypotheses and design, very little missing data, and sophisticated statistical methods. Finally, the medium to large effect sizes suggest that statistically significant results may be found with a larger, sufficiently powered, sample size.

The problem of child maltreatment and maternal substance abuse is undoubtedly a public health concern of the utmost significance (Magura & Laudet, 1996). Federal and state child welfare laws, most notably the federal Adoption and Safe Families Act (ASFA) of 1997, have resulted in dramatic changes in how dependency courts operate. Courts are no longer charged only with determining the validity of child abuse and neglect reports and then deciding whether to place the child in foster care. They must now also take a more active role in developing service plans for families and ensuring that each child is placed in a permanent and stable home (Hardin, 1996; Harrell & Goodman, 1999). The demands placed on child welfare systems and family courts resulting from the Adoption and Safe Families Act ASFA, combined with the growing number of substance-abusing parents involved in family court proceedings, have strained these systems' abilities to successfully resolve cases in the expeditious manner required by law.

Judicial and child welfare systems throughout the nation have turned to cooperative and non-adversarial drug courts as a setting where parents can acquire the tools needed to turn their lives around and become productive, drug-free members of society (Tauber & Snively, 1999). However, the lack of scientifically rigorous investigations of family drug courts severely limits public policy. Too many questions remain regarding the effectiveness, essential features, and influence of family drug courts on drug and non-drug outcomes. Even though family drug courts have an increasingly important role to play in promoting healthy families, few well-specified intervention programs aimed at drug-abusing mothers suitable for use in drug courts are available. There can be no doubt that further investigation into the processes and outcomes of dependency drug court is warranted, especially work that include the state-of-the-art methods incorporated in the current study plus a larger sample size, more comprehensive assessment of intervention fidelity, an investigation into the mechanisms of change, and perhaps even a benefit-cost analysis. Nevertheless, this pilot study adds to the field by suggesting that family drug courts, and perhaps especially ones that employ systematic family-based interventions such as the Engaging Moms Program, may be a viable approach to effectively addressing the problem of maternal substance abuse and child maltreatment. This study represents one step in the development of successful family drug court models, models that may have a significant impact on the health of the mother, the child, and the larger social problems associated with maternal drug addiction (Boles, Young, Moore, & DiPirro-Beard, 2007; Green, Furrer, Worcel, Burrus, & Finigan, 2007; Haack, Alemi, Nemes, & Cohen, 2004).

Acknowledgments

This article was supported by a grant from the National Institute on Drug Abuse, RO1 DA016733. From the Miami Dade County 11th District Circuit Court, we thank the Honorable Cindy Lederman and Sharon Abrams for facilitating this research, and the drug court counselors: Marlen Bouzon, Cynthia Howard, Amparo Carrera, Alexa Ortiz, and Tammy Rivers for their cooperation and participation in this study.

References

- Attkisson, CC.; Greenfield, TK. Client Satisfaction Questionnaire-8 and Service Satisfaction Scale-30. In: Mariush, ME., editor. *The use of psychological testing for treatment planning and outcome assessment*. Lawrence Erlbaum; Hillsdale, NJ: 1994. p. 402-420.
- Barth, RP.; Courtney, M.; Duerr Berrick, J.; Albert, V. *From child abuse to permanency planning: Child welfare services pathways and placements*. Aldine de Gruyter; Hawthorne, NY: 1994.
- Belenko S. *Research on drug courts: A critical review: 2001 update*. National Drug Court Institute Review. 2001
- Boles SM, Young NK, Moore T, DiPirro-Beard S. The Sacramento Dependency Drug Court, Development and outcomes. *Child Maltreatment* 2007;12:161–171. [PubMed: 17446569]

- Bureau of Justice Assistance (BJA). Drug Court Clearinghouse Project. Summary of Drug Court Activity by State and County. Jun 18. 2009 Available at: <http://www1.spa.american.edu/justice/documents/2153.pdf>
- Carise D, McLellan AT, Cacciola J, Love M, Cook T, Bovasso G, Lam V. Suggested specifications for a standardized Addiction Severity Index database. *Journal of Substance Abuse Treatment* 2001;20:239–244. [PubMed: 11516594]
- Catalano RF, Gainey RR, Fleming CB, Haggerty KP, Johnson NO. An experimental intervention with families of substance abusers: One-year follow-up of the focus on families project. *Addiction* 1999;94(2):241–254. [PubMed: 10396792]
- Choi S, Ryan JP. Completing substance abuse treatment in child welfare: The role of co-occurring problems and primary drug of choice. *Child Maltreatment* 2006;11:313–325. [PubMed: 17043316]
- Connell CM, Bergeron N, Katz KH, Saunders L, Tebes JK. Re-referral to child protective services: The influence of child, family, and case characteristics on risk status. *Child Abuse and Neglect* 2007;31:573–588. [PubMed: 17537504]
- Crawford AM, Pentz MA, Chou CP, Li C, Dwyer JH. Parallel developmental trajectories of sensation seeking and regular substance use in adolescents. *Psychology of Addictive Behaviors* 2003;17(3):179–192. [PubMed: 14498812]
- Curtis PA, McCullough C. The impact of alcohol and other drugs on the child welfare system. *Child Welfare* 1993;72(6):533–542. [PubMed: 8243114]
- Dakof GA, Cohen JB, Duarte E. Increasing family reunification for substance abusing mothers and their children: Comparing two drug court interventions. *Juvenile and Family Court Journal* 2009;60:11–23.
- Dakof GA, Quille TJ, Tejada MJ, Alberga LR, Bandstra E, Szapocznik J. Enrolling and retaining mothers of substance-exposed infants into drug abuse treatment. *Journal of Consulting and Clinical Psychology* 2003;71:764–772. [PubMed: 12924681]
- Delucchi K, Bostrom A. Small sample longitudinal clinical trial with missing data: A comparison of analytic methods. *Psychological Methods* 1999;4:158–172.
- Derogatis, LR. BSI: Administration, scoring and procedures manual- 3rd Ed.. National Computer Systems; Minneapolis, MN: 1993.
- Derogatis, LR.; Savitz, KL. The SCL-90-R, Brief Symptom Inventory, and Matching clinical rating scales. In: Maruish, ME., editor. *The use of psychological testing for treatment planning and outcome assessment-2nd Edition*. Lawrence Erlbaum Associates; Mahwah, NJ: 1999. p. 679-724.
- Dore MM, Doris JM, Wright P. Identifying substance abuse in maltreating families: A child welfare challenge. *Child Abuse and Neglect* 1995;19(5):531–543. [PubMed: 7664134]
- Drapela LA, Mosher C. The conditional effect of parental drug use on parental attachment and adolescent drug use: Social control and social development model perspectives. *Journal of Child and Adolescent Substance Abuse* 2007;16(3):63–87.
- Duncan, TE.; Duncan, SC.; Strycker, LA.; Li, F.; Alpert, A. *An introduction to latent variable growth curve modeling: Concepts, issues, and applications*. Lawrence Erlbaum; Mahwah, NJ: 1999.
- Edwards LP, Ray JA. Judicial perspectives on family drug treatment courts. *Juvenile and Family Court Journal* 2005:1–27.
- Feingold A. Effect sizes for growth-modeling analysis for controlled clinical trials in the same metric as for classical analysis. *Psychological Methods* 2009;14(1):43–53. [PubMed: 19271847]
- Gregoire KA, Schultz DJ. Substance-abusing child welfare parents: Treatment and child placement outcomes. *Child Welfare* 2001;80(4):433–452. [PubMed: 11480487]
- Grella CE, Needell B, Shi Y, Hser YI. Do drug treatment services predict reunification outcomes of mothers and their children in child welfare? *Journal of Substance Abuse Treatment* 2009;36(3):278–293. [PubMed: 18775623]
- Greenfield TK, Attkisson CC. Steps toward a multifactorial satisfaction scale for primary care and mental health services. *Evaluation and Program Planning* 1989;12:271–278.
- Green BL, Furrer C, Worcel S, Burrus S, Finnigan M. How effective are family drug courts? Outcomes from a four-site national study. *Child Maltreatment* 2007;12:43–59. [PubMed: 17218647]

- Green BL, Furrer C, Worcel S, Burrus S, Finnigan M. Building the evidence base for family drug treatment courts: Results from recent outcome studies. *Drug Court Review* 2009;6:53–82.
- Green BL, Rockhill A, Furrer C. Does substance abuse treatment make a difference for child welfare case outcomes? A statewide longitudinal analysis. *Children and Youth Services Review* 2007;29:460–473.
- Grogan-Kaylor A, Ruffolo MC, Ortega RM, Clarke J. Behaviors of youth involved in the child welfare system. *Child Abuse and Neglect* 2008;32:35–49. [PubMed: 18155294]
- Haack M, Alemi F, Nemes S, Cohen JB. Experience with family drug courts in three cities. *Substance Abuse* 2004;25:17–25. [PubMed: 16172089]
- Hardin M. Responsibilities and effectiveness of the juvenile court in handling dependency cases. *The Future of Children* 6 1996:111–125.
- Harrell, A.; Goodman, A. Review of specialized family drug courts: Key issues in handling child abuse and neglect cases. The Urban Institute; Washington, DC: 1999.
- Henggeler SW. Juvenile drug courts: emerging outcomes and key research issues. *Current Opinion in Psychiatry* 2007;20:242–246. [PubMed: 17415076]
- Horvath AO, Symonds BD. Relation between working alliance and outcome in psychotherapy: A meta-analysis. *Journal of Counseling Psychology* 1991;38(2):139–149.
- Huddleston, CW.; Marlowe, DB.; Casebolt, R. Painting the current picture: A national report card on drug courts and other problem-solving court programs in the United States. Vol. 21. National Drug Court Institute; Alexandria, VA: May. 2008
- Keller TE, Catalano RF, Haggerty KP, Fleming CB. Parent figure transitions and delinquency and drug use among early adolescent children of substance abusers. *American Journal of Drug and Alcohol Abuse* 2002;28(3):399–427. [PubMed: 12211358]
- Kelley ML, Fals-Stewart W. Psychiatric disorders of children living with drug-abusing, alcohol-abusing, and non-substance-abusing fathers. *Journal of the American Academy of Child and Adolescent Psychiatry* 2004;43:621–628. [PubMed: 15100569]
- Kelley ML, Fals-Stewart W. Treating paternal drug abuse using Learning Sobriety Together: Effects on adolescents versus children. *Drug and Alcohol Dependence* 2008;92:228–238. [PubMed: 17890017]
- Kivlighan DM Jr. Shaughnessy P. An analysis of the development of the working alliance using hierarchical linear modeling. *Journal of Counseling Psychology* 1995;42:338–349.
- Liddle, HA.; Dakof, GA.; Diamond, GS. Adolescents and substance abuse: Multidimensional family therapy in action. In: Kaufman, E.; Kaufman, P., editors. Family therapy approaches with drug and alcohol problems. 2nd edition. Gardner; 1991. p. 120-171.
- Liddle HA, Rowe CL, Dakof GA, Henderson CE, Greenbaum PE. Early intervention for adolescent substance abuse: 12-month treatment outcomes of family-based versus group treatment. *Journal of Consulting and Clinical Psychology* 2009;77:12–25. [PubMed: 19170450]
- Little, RJA.; Rubin, DB. Statistical analysis with missing data. Wiley; New York: 1987.
- Locke TF, Newcomb MD. Child maltreatment, parent alcohol- and drug-related problems, polydrug problems, and parenting practices: A test of gender differences and four theoretical perspectives. *Journal of Family Psychology* 2004;18(1):120–134. [PubMed: 14992615]
- Magura S, Laudet AB. Parental substance abuse and child maltreatment: Review and implications for interventions. *Children and Youth Services Review* 1996;18:193–220.
- Marcenko MO, Kemp SP, Larson NC. Childhood experiences of abuse, later substance use, and parenting outcomes among low-income mothers. *American Journal of Orthopsychiatry* 2000;70:316–326. [PubMed: 10953778]
- McLellan AT, Alterman AI, Cacciola J, Metzger D, O'Brien CP. A new measure of substance abuse treatment: Initial studies of the Treatment Services Review. *Journal of Nervous and Mental Disease* 1992;180:101–110. [PubMed: 1737971]
- Miller, WR.; Tonigan, JS. Psychiatry and Addictive Behavior. Vol. 10. 1996. Assessing drinkers' motivation for change: The stages of change readiness and treatment eagerness scale (SOCRATES); p. 81-89.
- Monchick, R.; Scheyett, A.; Pfeifer, J. Drug Court Case Management: Role, Function, and Utility. National Drug Court Institute; Alexandria, VA: 2006.

- Muthén, L.; Muthén, B. Mplus user's guide [Computer software]. Muthén & Muthén; Los Angeles, CA: 19982009.
- National Associate of Drug Court Professionals (NADCP). Defining drug courts: The key components. Department of Justice, Drug Courts Program Office; Washington DC: 1997. Retrived from the Web: [www.ndci.org/sites/default/files/ndci/Key Components.pdf](http://www.ndci.org/sites/default/files/ndci/Key%20Components.pdf)
- Undersma SJ, Chaffin M, Simpson S, LeBreton J. The Brief Child Abuse Potential inventory: Development and validation. *Journal of Clinical Child and Adolescent Psychology* 2005;34:301–311. [PubMed: 15901230]
- Rounsaville BJ, Carroll KM, Onken LS. A stage model of behavioral therapies research: getting started and moving on from stage I. *Clinical Psychology Science and Practice* 2001;8(2):133–142.
- Pach NM. An overview of operational family dependency treatment courts. *Drug Court Review* 2009;VI: 67–110.
- Parr, J.; Greenfield, TK. Methadone maintenance treatment at Cairns Base Hospital: client characteristics and client satisfaction. Cairns District Health Service, Alcohol, Tobacco & Other Drugs Service; Cairns, Queensland, Australia: 2000.
- Semidei J, Radel LF, Nolan C. Substance abuse and child welfare: Clear linkages and promising responses. *Child Welfare* 2001;80:109–128. [PubMed: 11291897]
- Stanger C, Higgins ST, Bickel WK, Elk R, Grabowski J, Schmitz J, et al. Behavioral and emotional problems among children of cocaine- and opiate-dependent parents. *Journal of the American Academy of Child and Adolescent Psychiatry* 1999;38(4):421–428. [PubMed: 10199114]
- Strauss MA. Measuring intrafamily conflict and violence: The conflict tactics scale. *Journal of Marriage and Family* 1979;41:75–87.
- Tauber, JJ.; Snavelly, KR. Drug courts: a research agenda. National Drug Court Institute; Alexandria, VA: 1999.
- Walsh C, MacMillan HL, Jamieson E. The relationship between parental substance abuse and child maltreatment, Findings from the Ontario Health Supplement. *Child Abuse and Neglect* 2003;27:1409–1425. [PubMed: 14644058]
- Wilson DB, Mitchell O, Mackenzie DL. A systematic review of drug court effects on recidivism. *Journal of Experimental Criminology* 2006;2:459–487.
- Wolock I, Magura S. Parental substance abuse as a predictor of child maltreatment re-reports. *Child Abuse and Neglect* 1996;20:1183–1193. [PubMed: 8985609]
- Worcel SD, Furrer CJ, Green BL, Burrus SWM, Finigan MW. Effects of family treatment drug courts on substance abuse and child outcomes. *Child Abuse Review* 2008;17:427–443.

Table 1

Sample Characteristics

Variable	EMP	ICMS	Overall
Age	29.1 (7.6)	31.2 (14.0)	30.2 (11.4)
Ethnicity [<i>n</i> (%)]			
Black	14 (45)	12 (39)	26 (42)
Hispanic	11 (36)	11 (35)	22 (35)
White	6 (19)	8 (26)	14 (23)
Education [<i>n</i> (%)]			
< High School Graduation	16 (52)	19 (61)	35 (57)
Graduated High School/GED	15 (48)	8 (26)	23 (37)
Some College	0 (0)	4 (13)	4 (6)
Employment Status [<i>n</i> (%)]			
Employed	8 (26)	10 (32)	18 (29)
Unemployed	23 (74)	21 (68)	44 (71)
Monthly Family Income [<i>M</i> (<i>SD</i>)]	674 (652)	1016 (1138)	845 (939)
Median	507	561	561
Number of Children [<i>M</i> (<i>SD</i>)]	2.5 (1.6)	2.5 (1.4)	2.5 (1.5)
Median	2.0	3.0	2.0
Marital Status [<i>n</i> (%)]			
Married	2 (6)	4 (13)	6 (10)
Divorced/Separated	7 (23)	8 (26)	15 (24)
Never Married	22 (71)	19 (61)	41 (66)
Variable	EMP	ICMS	Overall
% with Physical Abuse Hx	48	61	55
% with Sexual Abuse Hx	39	32	36
Age at First Use [<i>M</i> (<i>SD</i>)]	17.3 (2.8)	16.1 (4.5)	16.7 (3.7)
Drug of Choice [<i>n</i> (%)]			
Alcohol	1 (3)	2 (7)	3 (5)
Cannabis	3 (10)	3 (10)	6 (10)
Cocaine	4 (13)	6 (19)	10 (16)
Other Sedatives	0 (0)	1 (3)	1 (2)
Alcohol and Polydrug use	17 (55)	13 (42)	30 (48)
Polydrug use but no alcohol	6 (19)	6 (19)	12 (19)
Mental Health History [<i>n</i> (%)]			
Serious Depression	23 (68)	21 (68)	44 (68)
Serious Anxiety	17 (55)	17 (55)	34 (55)
Hallucinations	4 (13)	4 (13)	8 (13)
Suicidal Ideation	8 (26)	4 (13)	12 (19)
Total Lifetime Arrests [<i>M</i> (<i>SD</i>)]	2.3 (4.1)	3.9 (12.5)	3.1 (9.3)

Note. M = Mean, SD = Standard Deviation.

Table 2

Total hours of intervention contact with mother alone, with family, and with extrafamilial domains by intervention condition

Variable	EMP	ICMS
Contact Hours [<i>m (SD)</i>]		
Mom	25.62 (12.91)	25.3(13.68)
Median	26.98	29.05
Family	6.92 (9.46)**	3.83 (3.92)**
Median	3.08	2.08
Mom-Extrafamilial	5.95 (7.03)	8.23 (5.28)
Median	3	8
Total Contacts	38.48 (20.26)	37.26 (19.15)
Median	36.61	42.45

* = $p < .05$,

** = $p < .01$,

*** = $p < .001$

Table 3

Frequency and percentage of EMP and ICMS child welfare dispositions at 18 months after enrollment in drug court

Disposition [<i>n</i> (%)]			
	EMP	ICMS	Overall
TPR, Placed in Foster Care	4 (13)	9 (29)	13 (21)
TPR, Placed w/Relatives	3 (10)	5 (16)	8 (13)
No TPR, Placed w/ Relatives	6 (19)	3 (10)	9 (14)
Joint Custody	2 (6)	2 (6)	4 (6)
Sole Custody	16 (52)	12 (39)	28 (46)

Note. TPR = Terminated Parental Rights

Table 4
Observed Sample Means and Standard Deviations for Secondary Outcome Variables

Outcome Measure	Intake	3 Month	6 Month	9 Month	12 Month	18 Month
	<i>M (SD)</i>					
ASI Alcohol Scale						
EMP	.21 (.26)	.02 (.07)	.02 (.08)	.03 (.11)	.03 (.09)	.00 (.01)
ICMS	.11 (.16)	.02 (.08)	.03 (.12)	.02 (.04)	.01 (.02)	.02 (.05)
ASI Drug Scale						
EMP	.14 (.14)	.02 (.04)	.04 (.09)	.02 (.06)	.04 (.07)	.02 (.07)
ICMS	.13 (.14)	.03 (.06)	.02 (.08)	.02 (.08)	.02 (.06)	.03 (.05)
ASI Family/Social Scale						
EMP	.29 (.27)	.05 (.10)	.05 (.09)	.05 (.08)	.07 (.11)	.02 (.06)
ICMS	.22 (.25)	.09 (.16)	.06 (.11)	.05 (.07)	.03 (.08)	.03 (.06)
ASI Medical Scale EMP						
EMP	.15 (.29)	.09 (.25)	.14 (.27)	.04 (.13)	.03 (.09)	.02 (.11)
ICMS	.21 (.34)	.10 (.27)	.06 (.23)	.05 (.21)	.06 (.19)	.11 (.25)
ASI Employment Scale						
EMP	.75 (.28)	.82 (.24)	.80 (.19)	.65 (.31)	.62 (.32)	.65 (.31)
ICMS	.77 (.25)	.82 (.25)	.77 (.26)	.61 (.33)	.62 (.26)	.63 (.32)
ASI Mental Health Scale						
EMP	.31 (.27)	.11 (.20)	.09 (.13)	.06 (.09)	.09 (.12)	.06 (.11)
ICMS	.27 (.21)	.12 (.18)	.12 (.17)	.04 (.22)	.11 (.14)	.09 (.21)
BCAP Risk Scale						
EMP	12.42 (5.77)	8.14 (4.98)	7.90 (5.74)	6.46 (4.79)	7.31 (6.43)	4.81 (3.85)
ICMS	11.03 (5.40)	9.12 (5.83)	7.36 (5.77)	6.48 (5.23)	7.52 (5.19)	4.83 (3.84)
BSI Global Severity Index						
EMP	.91 (.92)	.30 (.42)	.32 (.60)	.08 (.15)	.24 (.52)	.16 (.35)
ICMS	.74 (.68)	.53 (.60)	.25 (.43)	.20 (.30)	.28 (.32)	.17 (.36)
CTS Physical Violence						
EMP	2.28 (11.13)	3.93 (20.79)	1.07 (4.84)	1.14 (5.33)	.00 (.00)	.00 (.00)
ICMS	3.57 (19.54)	1.41 (5.92)	.15 (.78)	.00 (.00)	.07 (.38)	.14 (.76)
Positive Urine Screen [n (%)]						

Outcome Measure	Intake	3 Month		6 Month		9 Month		12 Month		18 Month	
	<i>M (SD)</i>										
EMP	12 (38.7)	5 (16.1)	4 (12.9)	1 (3.2)	8 (25.8)	2 (6.5)	7 (22.6)	5 (16.1)	2 (6.5)		
ICMS	10 (32.3)	2 (6.5)	5 (16.1)	6 (19.4)							

Note. ASI = Addiction Severity Index, BCAP = Brief Child Abuse Potential, BSI = Brief Symptom Inventory, CTS = Conflict Tactics Scale

Table 5
Tests of slope main effects and growth factor by treatment interactions for outcome variables.

Outcome Measure	Growth Factor Mean Intake - 3 months			Growth Factor Mean 3-18 months			Treatment Differences Intake - 3 months			Treatment Differences 3-18 months				
	Intercept Estimate	SE	Slope Estimate	SE	Slope Estimate	SE	Intercept Estimate	SE	Slope Estimate	SE	Intercept Estimate	SE	Slope Estimate	SE
ASI Alcohol	0.16***	0.03	-0.14***	0.03	<0.01	<0.01	-0.10	0.05	0.09	0.05	0.01	0.05	<0.01	<0.01
ASI Drug	0.13***	0.02	-0.11***	0.02	<0.01	<0.01	-0.01	0.04	0.01	0.04	<0.01	0.04	<0.01	0.01
ASI Family/Social	0.25***	0.03	-0.18***	0.03	-0.01	<0.01	-0.08	0.07	0.08	0.07	<0.01	0.07	<0.01	0.01
ASI Medical	0.18***	0.04	-0.09*	0.04	-0.01	0.01	0.06	0.08	-0.10	0.09	0.02	0.09	0.02	0.01*
ASI Employment	0.76***	0.03	0.05*	0.02	-0.05***	0.01	0.02	0.07	-0.03	0.05	<0.01	0.05	<0.01	<0.01
ASI Mental Health	0.29***	0.03	-0.18***	0.03	-0.01	0.01	-0.03	0.06	0.05	0.07	<0.01	0.07	<0.01	0.01
BCAP Risk	11.77***	0.71	-3.20***	0.65	-0.84***	0.19	-1.32	1.40	1.77	1.23	-0.17	1.23	-0.17	0.39
BSI GSI	0.83***	0.10	-0.51***	0.10	-0.04**	0.02	-0.17	0.20	0.34	0.19	-0.04	0.19	-0.04	0.03
CTS NVD	3.93***	0.97	-0.62	1.30	1.70***	0.51	-1.27	1.96	3.31	2.52	-1.21	2.52	-1.21	0.87
Urinalysis	N/A	N/A	-1.54**	0.48	-0.51	0.47	-0.32	0.59	-0.20	0.90	0.40	0.90	0.40	0.55

Note. ASI = Addiction Severity Index, BCAP = Brief Child Abuse Potential, BSI GSI = Brief Symptom Inventory Global Severity Index, CTS = Conflict Tactics Scale Nonviolent Discipline

* $p < .05$.

** $p < .01$

*** $p < .001$.